

MAS114 Problems

Sheet 11 (Week 12)

Preamble

Themes from this week (ask your tutorial staff if you're stuck): 1. The Sandwich Lemma. 2. General facts about convergent sequences. 3. Cauchy sequences. 4. Constructing the real numbers.

Work on these problems one at a time in small groups (of around four). For many problems there is designed to be a lot of work that's best shared; and for others discussion is vital to understanding.

1

In this question, we'll write $f(x) \prec g(x)$ to mean "there is an X such that $f(x) < g(x)$ for all $x > X$ ", which might be thought of informally as saying "when x gets large, eventually we always have $f(x) < g(x)$ ". So, for example, $x^4 \prec x^5$, as we have $x^4 < x^5$ for all $x > 1$.

Note that if $f(x) \prec g(x)$, and $g(x) \prec h(x)$, then $f(x) \prec h(x)$.

- (i) For each of the three \prec symbols in the following chain, find an X such that the inequality holds for all $x > X$:

$$1000000x \ln x \prec 1000x^2 \prec x^{2.001} \prec 1.000001^x.$$

- (ii) Arrange the following functions to form a chain ordered by \prec like the one above:

$$\bullet (\ln x)^{\sqrt{x}} \quad \bullet (\sqrt{x})^{\ln x} \quad \bullet \frac{1000x^2}{\ln x} \quad \bullet \frac{e^{\sqrt{x}}}{1000000} \quad \bullet \frac{x^2}{\ln(\ln x)}$$

In each of the four places you use it, prove that the relation \prec holds.

For discussion: Can you prove that your chain is correctly ordered by \prec ?

Given any two functions $f(x)$ and $g(x)$, must we have $f(x) \prec g(x)$, or $g(x) \prec f(x)$, or $f(x) = g(x)$?

2

Note that the statement “the sequence a_0, a_1, \dots does not converge to x ” means “there exists some ϵ , such that for all N there exists $n > N$ such that $|a_n - x| \geq \epsilon$ ”. Write down a careful proof, using that, that the sequence defined by $a_n = n$ does not converge to any x .

For discussion: Does it feel easier or harder to prove things *do not* converge than that they do? Is this reasonable, or just because you’re not used to it?

3

(i) Prove that $2^n \geq n^2$ for all integers $n \geq 4$.

(ii) Prove that we have

$$0 \leq \frac{n}{2^n} \leq \frac{1}{n}$$

for all n .

(iii) Conclude using the Sandwich Lemma that the sequence $\frac{n}{2^n}$ converges to zero.

4

Given a sequence a_1, a_2, a_3, \dots , a *subsequence* is something formed by choosing some of the terms only. For example, $a_2, a_3, a_5, a_7, \dots$ is a subsequence. Can you find a sequence a_1, a_2, a_3, \dots with the property that, for any real number $0 \leq x \leq 1$, there is a subsequence of a_1, a_2, a_3, \dots which converges to x ?