

# MAS114: Lecture 3

James Cranch

<http://cranch.staff.shef.ac.uk/mas114/>

2017–2018

# Announcement: exercises

## Announcement: exercises

Have you found the homework online yet?

## Announcement: exercises

Have you found the homework online yet?

You need to hand it in at the beginning of your problems class at the end of the week.

# Unions

# Unions

Let  $A$  and  $B$  be sets. We define their *union*  $A \cup B$  to contain exactly the things that are in one set or the other (or both):

$$A \cup B = \{x \mid x \in A \text{ or } x \in B\}.$$

# Unions

Let  $A$  and  $B$  be sets. We define their *union*  $A \cup B$  to contain exactly the things that are in one set or the other (or both):

$$A \cup B = \{x \mid x \in A \text{ or } x \in B\}.$$

That notation is called a *set comprehension*: the thing on the left of the vertical bar are the things we want to put in the set, and the things on the right of the vertical bar are the conditions under which we put them in. We'll use them a lot.

# Intersections



# Intersections

Similarly, we define the *intersection*  $A \cap B$  to contain exactly the things that are in both sets:

$$A \cap B = \{x \mid x \in A \text{ and } x \in B\}.$$

# Differences

# Differences

Lastly, we define the *difference*  $A \setminus B$  to be the things which are in  $A$  but not in  $B$ :

$$A \setminus B = \{x \mid x \in A \text{ and } x \notin B\}.$$

# Equality of sets

# Equality of sets

Two sets  $A$  and  $B$  are equal if they have the same members.

# Equality of sets

Two sets  $A$  and  $B$  are equal if they have the same members.  
A straightforward way of proving this is often to show that  $A \subset B$   
and  $B \subset A$ .

# Equality of sets

Two sets  $A$  and  $B$  are equal if they have the same members. A straightforward way of proving this is often to show that  $A \subset B$  and  $B \subset A$ . That is, in words, two sets  $A$  and  $B$  are equal if every element of  $A$  is an element of  $B$  and every element of  $B$  is an element of  $A$ .

# An example



## An example

Here's an example of this proof strategy:

## An example

Here's an example of this proof strategy:

### Proposition

*Let  $A$ ,  $B$  and  $C$  be three sets. We have*

$$A \cap (B \cup C) = (A \cap B) \cup (A \cap C).$$

## An example

Here's an example of this proof strategy:

### Proposition

*Let  $A$ ,  $B$  and  $C$  be three sets. We have*

$$A \cap (B \cup C) = (A \cap B) \cup (A \cap C).$$

Proof.



# Some comments

## Some comments

That was the first example of a formal proof in this course.

## Some comments

That was the first example of a formal proof in this course. You'll have to write many proofs like this yourself, in assessed homework and in the exam.

## Some comments

That was the first example of a formal proof in this course. You'll have to write many proofs like this yourself, in assessed homework and in the exam. Though we'll discuss it in depth later, it may be worth observing the style from the beginning.

## Some comments

That was the first example of a formal proof in this course. You'll have to write many proofs like this yourself, in assessed homework and in the exam. Though we'll discuss it in depth later, it may be worth observing the style from the beginning. One big mistake that many beginner mathematicians make is *not using words to explain the flow of the argument*.



# A warning

## A warning

What we are practising here is called *naïve set theory*. What's so naïve about it?

## A warning

What we are practising here is called *naïve set theory*. What's so naïve about it?

The Welsh mathematician Bertrand Russell realised in 1901 that there are serious problems with being allowed to form sets carelessly:

## A warning

What we are practising here is called *naïve set theory*. What's so naïve about it?

The Welsh mathematician Bertrand Russell realised in 1901 that there are serious problems with being allowed to form sets carelessly:

### Paradox

*Suppose there is a set  $S$  of all sets which are not elements of themselves:*

$$S = \{A \mid A \notin A\}.$$

*This creates a contradiction.*

## A warning

What we are practising here is called *naïve set theory*. What's so naïve about it?

The Welsh mathematician Bertrand Russell realised in 1901 that there are serious problems with being allowed to form sets carelessly:

### Paradox

*Suppose there is a set  $S$  of all sets which are not elements of themselves:*

$$S = \{A \mid A \notin A\}.$$

*This creates a contradiction.*

Proof.



# Modern set theory

# Modern set theory

As a result of this paradox, modern set theorists impose strict rules on what sets can be formed, with the aim of banning this particular beast and everything like it.

# Modern set theory

As a result of this paradox, modern set theorists impose strict rules on what sets can be formed, with the aim of banning this particular beast and everything like it. However, you probably won't need to worry about this, unless you take a course in set theory later in your mathematical careers.



# Functions

# Functions

A function is to be thought of as a machine that takes an element of one set and gives you an element of another.

# Functions

A function is to be thought of as a machine that takes an element of one set and gives you an element of another. Here's a formal definition:

# Functions

A function is to be thought of as a machine that takes an element of one set and gives you an element of another. Here's a formal definition:

## Definition

Given sets  $A$  and  $B$ , a *function* (sometimes called a *map*)  $f : A \rightarrow B$  gives for each element  $a \in A$  a unique element  $f(a) \in B$ .

# Examples of functions

# Examples of functions

Examples include:

# Examples of functions

Examples include:

- ▶ the function  $f : \mathbb{Z} \rightarrow \mathbb{Z}$  defined by  $f(n) = n^2 - 7$ .

# Examples of functions

Examples include:

- ▶ the function  $f : \mathbb{Z} \rightarrow \mathbb{Z}$  defined by  $f(n) = n^2 - 7$ .
- ▶ the function  $g : \mathbb{Q} \rightarrow \{4, 6\}$  defined by

$$g(x) = \begin{cases} 4, & \text{if } x = 3/7 \text{ or } x = -14/17; \\ 6, & \text{otherwise.} \end{cases}$$



# Domain and codomain

# Domain and codomain

The set  $A$  is called the *domain* of  $f$ , and  $B$  is called the *codomain* of  $f$ .

## Domain and codomain

The set  $A$  is called the *domain* of  $f$ , and  $B$  is called the *codomain* of  $f$ . We call  $f(a)$  the *value* of  $f$  at  $a$ , or the *image* of  $a$  under  $f$ .

# Domain / Image / Range

## Domain / Image / Range

Consider the “age in years” function from the set of people in this room to the natural numbers.

## Domain / Image / Range

Consider the “age in years” function from the set of people in this room to the natural numbers.

The **domain** of this function is the set of values you’re permitted to apply it to.

## Domain / Image / Range

Consider the “age in years” function from the set of people in this room to the natural numbers.

The **domain** of this function is the set of values you’re permitted to apply it to. This is the set of people in this room, because I said so.

## Domain / Image / Range

Consider the “age in years” function from the set of people in this room to the natural numbers.

The **domain** of this function is the set of values you’re permitted to apply it to. This is the set of people in this room, because I said so.

The **codomain** of this function is the set of values it is *permitted* to take.



## Domain / Image / Range

Consider the “age in years” function from the set of people in this room to the natural numbers.

The **domain** of this function is the set of values you’re permitted to apply it to. This is the set of people in this room, because I said so.

The **codomain** of this function is the set of values it is *permitted* to take. This is the set  $\mathbb{N}$  of natural numbers, because I said so.

## Domain / Image / Range

Consider the “age in years” function from the set of people in this room to the natural numbers.

The **domain** of this function is the set of values you’re permitted to apply it to. This is the set of people in this room, because I said so.

The **codomain** of this function is the set of values it is *permitted* to take. This is the set  $\mathbb{N}$  of natural numbers, because I said so.

Some people like to talk about the **image** of this function, being the set of values it actually takes in practice.

## Domain / Image / Range

Consider the “age in years” function from the set of people in this room to the natural numbers.

The **domain** of this function is the set of values you’re permitted to apply it to. This is the set of people in this room, because I said so.

The **codomain** of this function is the set of values it is *permitted* to take. This is the set  $\mathbb{N}$  of natural numbers, because I said so.

Some people like to talk about the **image** of this function, being the set of values it actually takes in practice. This might (perhaps) be the set

$$\{18, 19, 20, 34\}.$$

## Domain / Image / Range

Consider the “age in years” function from the set of people in this room to the natural numbers.

The **domain** of this function is the set of values you’re permitted to apply it to. This is the set of people in this room, because I said so.

The **codomain** of this function is the set of values it is *permitted* to take. This is the set  $\mathbb{N}$  of natural numbers, because I said so.

Some people like to talk about the **image** of this function, being the set of values it actually takes in practice. This might (perhaps) be the set

$$\{18, 19, 20, 34\}.$$

The **range** is not a phrase that’s used consistently:

## Domain / Image / Range

Consider the “age in years” function from the set of people in this room to the natural numbers.

The **domain** of this function is the set of values you’re permitted to apply it to. This is the set of people in this room, because I said so.

The **codomain** of this function is the set of values it is *permitted* to take. This is the set  $\mathbb{N}$  of natural numbers, because I said so.

Some people like to talk about the **image** of this function, being the set of values it actually takes in practice. This might (perhaps) be the set

$$\{18, 19, 20, 34\}.$$

The **range** is not a phrase that’s used consistently:

- ▶ some people use it to mean the codomain;

## Domain / Image / Range

Consider the “age in years” function from the set of people in this room to the natural numbers.

The **domain** of this function is the set of values you’re permitted to apply it to. This is the set of people in this room, because I said so.

The **codomain** of this function is the set of values it is *permitted* to take. This is the set  $\mathbb{N}$  of natural numbers, because I said so.

Some people like to talk about the **image** of this function, being the set of values it actually takes in practice. This might (perhaps) be the set

$$\{18, 19, 20, 34\}.$$

The **range** is not a phrase that’s used consistently:

- ▶ some people use it to mean the codomain;
- ▶ some people use it to mean the image;

## Domain / Image / Range

Consider the “age in years” function from the set of people in this room to the natural numbers.

The **domain** of this function is the set of values you’re permitted to apply it to. This is the set of people in this room, because I said so.

The **codomain** of this function is the set of values it is *permitted* to take. This is the set  $\mathbb{N}$  of natural numbers, because I said so.

Some people like to talk about the **image** of this function, being the set of values it actually takes in practice. This might (perhaps) be the set

$$\{18, 19, 20, 34\}.$$

The **range** is not a phrase that’s used consistently:

- ▶ some people use it to mean the codomain;
- ▶ some people use it to mean the image;
- ▶ some (confused) people, who don’t know the difference, use it inconsistently to mean both.